Iranian Journal of Animal Biosystematics (IJAB)

Vol.15, No.1, 107-116, 2019

ISSN: 1735-434X (print); 2423-4222 (online)

DOI: 10.22067/ijab.v15i1.49697

# Hoverfly fauna (Diptera: Syrphidae) of Torbate Heydarie with a new record for the fauna of Razavi Khorasan province

Shojaei Hesari, E. and Pashaei Rad, Sh.\*

Department of Zoology, Faculty of Biological Science, G.C.Shahid Beheshti University, Tehran, Iran

(Received: 8 September 2018; Accepted: 14 May 2019)

The present study, during 2010-2011 was carried out in Torbat-e Heydarie (Razavi Khorasan Province). Adult speciemens were collected randomly with butterfly net. A totally, 28 species belong to 14 genus, two subfamilies were identified. All species are recorded for the first time from the studied area.

Key words: Hoverfly, Khorasan province, fauna, Iran, Diptera

#### INTRODUCTION

The hover flies or flower flies (Diptera: Syrphidae) are common in all areas from the Arctic to the Antarctic (Rotheray & Gilbert, 1999). About 6000 species in 200 genera and 40 tribes have been described (Thompson & Rotheray, 1998), and has been divided into three subfamilies: Microdontinae, Eristalinae and Syrphinae (Sommagio, 1999).

The adults mainly feed on nectar and pollen. The females must consume pollen since they need the proteins and amino acids of the pollen for maturation of their eggs. Nectar has only a small amount of these substances but much sugar. It is the fuel for the flies, enabling them to fly and hover actively. Being regular visitors of flowers, hoverflies are important pollinators of various plants including vegetables and fruit trees (Ghahari et al., 2008)

In addition, the larva of numerous species of hover flies, particularly subfamily of Syrphinae, are one of the most important predators of aphids and other pests which play important roles in their biological control (Gilbert, 1981). The Iranian syrphid fauna contain almost 188 species belong to 52 genera has been studied by some taxonomists: Farahbakhsh, 1961; Kuznetzov, 1985; Peck, 1988; Modarres awal, 1994; Melkeshi *et al.*, 1998; Gol Mohammad Zadeh Khiaban, 2000; Dousti, 2000; Gharali, 2000; Poor Rabi, 2000; Musavian, 2001; Sadeghi, 2001; Azarkhsh, 2001; Poor Ghasem, 2001; Keyvanfar & Sadeghi, 2002; Goldasteh, 2002; Sadeghi, 2003; Sadeghi, Kayvanfar & Jajvandan, 2006; Amiri Moghadam, 2004; Gilasian, 2005; Dousti *et al.*, 2006; Mehrabi *et al.*, 2006; Kayvanfar & Bagherian, 2006; Hosseini & Sadeghi, 2007; Ghahari *et al.*, 2008; Ashraphi & Pashaei Rad, 2010; Khaghaninia, 2010; Ehteshamnia *et al.*, 2010; Jalilian *et al.*, 2010; Gilasian & Sorokina, 2011; Ahmadian & Pashaei Rad, 2012; Kazerani *et al.*, 2012; Vosughian *et al.*, 2013; Khaghaninia & Hosseini, 2013; Kazerani *et al.*, 2014; Jalilian, 2014.

The Syrphid fauna of Razavi Khorasan province contain almost 39 species belong to 17 genera has been studied by: Musavian, 2001; Sadeghi, 2001; Keyvanfar & Sadeghi, 2002; Sadeghi & Keyvanfar, 2002; Sadeghi, 2003; Amiri Moghadam, 2004; Sadeghi, Kayvanfar & Jajvandian, 2006; Kayvanfar & Bagherian, 2006; Hosseini & Sadeghi, 2007. Torbat-e Heydarie city (35°16'N 59°14'E) is located in the center of Razavi Khorasan Province (Fig. 1) (Nami, 2010).




FIGURE 1. Torbat-e Heydarie city (35°16'N 59°14'E) in Razavi Khorasan province.

# Material and Methods

Three sampling localities include: Hesar, Robat Sang and Safi Abad have been shown in Table1. Species collection is done in four durations: 5-16 May 2010, 11-22 July 2010, 4-16 September 2010 and 5-16 May 2011. In this research, species are collected by a butterfly net with two random and sporadic ways. The best time for species collection is 7-10 am and 4-7 pm.

The collected insects were killed using killing jar then transferred to the laboratory, were pinned and were identified by valid keys such as Vockeroth & Thompson (1981), Bei- Bienko (1988), Stubbs & Falk (2002), Sorokina (2009), Speight (2010). Identifications were confirmed by Dr. Barkalov from Institute of Animal Systematics and Ecology (Russia).

#### **RESULTS**

A total of 526 specimens belong to 28 species, 14 genera and two subfamilies were identified, of which nine genera belong to the subfamily Syrphinae and five genera belong to subfamily Eristalinae. All species are recorded for the first time from Torbat-e Heydarie city. *Neoascia oblique* (Coe, 1940) was a new record for the fauna of Razavi Khorasan province.

# **SPECIES LIST**

#### Subfamily Syrphinae

# Chrysotoxum bicinctum Linnaeus, 1758

Material examined: 5 specimens  $(2 \circlearrowleft \circlearrowleft, 3 \hookrightarrow \circlearrowleft)$ , Hesar, 5-16 May 2010 (2 specimens), 4-16 September 2010 (3 specimens), *leg.* Elahe Shojaei.

Flowers visited: white umbellifers; *Achillea ptarmica*, *Alisma plantago-aquatica*, *Carduus* sp., *Crataegus* sp., *Hypochoeris* sp., *Potentilla erecta*, *Ranunculus* sp., *Rubus* sp. (Speight, 2011).

Distribution in Iran: Razavi Khorasan (Sadeghi, Kayvanfar & Jajvandan, 2006).

# Episyrphus balteatus De Geer, 1776

Material examined: 102 specimens ( $43 \stackrel{?}{\circlearrowleft} \stackrel{?}{\circlearrowleft}$ ,  $59 \stackrel{?}{\hookrightarrow} \stackrel{?}{\hookrightarrow}$ ), Hesar (67 specimens), Robat Sang (20 specimens), Safi Abad (15 specimens), 5-16 May 2010 (38 specimens), 11-22 July 2010 (17 specimens), 4-16 September 2010 (25 specimens), 5-16 May 2011 (22 specimens), leg. Elahe Shojaei.

Flowers visited: visits a wide range of white and yellow flowers, from trees to low-growing plants and including nectar less flowers; also visits pink flowers such as *Cirsium* sp. and *Succisa* sp.. In October/ November the species uses flowers of plants such as *Arbutus unedo*, *Hedera* sp. and *Viburnum tinus* (Speight, 2011).

Distribution in Iran: Ardebil, East Azerbayjan, Fars, Gilan, Golestan, Gorgan, Hamedan, Ilam, Isphahan, Kerman, Khuzestan, Lorestan, Maraghe, Mazandaran, Qazvin, Razavi Khorasan, Sistan and Baluchistan, West Azerbayjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Jalilian et al., 2010; Vosughian et al., 2013).

#### Eupeodes corollae Fabricius, 1794

Material examined: 83 specimens ( $35 \circlearrowleft \circlearrowleft$ ,  $48 \hookrightarrow \circlearrowleft$ ), Hesar (40 specimens), Robat Sang (26 specimens), Safi Abad (17 specimens), 5-16 May 2010 (19 specimens), 11-22 July 2010 (10 specimens), 4-16 September 2010 (17 specimens), 5-16 May 2011 (37 specimens), leg. Elahe Shojaei.

Flowers visited: Umbellifers; Achillea millefolium, Campanula rapunculoides, Chrysanthemum sp., Cirsium sp., Eschscholzia californica, Galeopsis sp., Hypericum sp., Leontodon sp., Origanum vulgare, Potentilla erecta, Ranunculus sp., Rubus fruticosus, Salix sp., Senecio sp., Tripleurospermum inodoratum, Tussilago sp. (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Fars, Golestan, Gorgan, Hamedan, Ilam, Isphahan, Khuzestan, Maraghe, Mazandaran, North Khorasan, Qazvin, Razavi Khorasan, Semnan, Sistan and Baluchistan, Tehran, West Azerbayjan, Zanjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Jalilian *et al.*, 2010; Vosughian *et al.*, 2013).

<b>75</b>		0	1.		•
TARLE 1		Samo	elmo	station	properties.
	•	Janip	cmig	Station	properties.

Number of station	Station	geographic coordinate	Mean Temprature	Mean Humidity	Plant species	Specimens collected
1	Hesar	35°41 <sup>′</sup> N 58°80 <sup>′</sup> E	24°C	41.5	Astragalus cancellatus Gundelia tournefortii Descurainia Sophia Convolvolus sp. Acanthophyllum sp.	272
2	Robat Sang	35°55′N 59°20′E	25°C	34	Convolvolus sp. Mentha sp. Poacea Trifolium sp. Medicayo sp.	132
3	Safi Abad	35°29 <sup>°</sup> N 59°63 <sup>°</sup> E	35°C	28	Convolvolus sp. Alhagi sp. Oryza sp. Prosopis farcta	122

# Eupeodes latifasciatus Macquart, 1829

**Material examined:** 7 specimens ( $2 \circlearrowleft \circlearrowleft$ ,  $5 \hookrightarrow \hookrightarrow$ ), Hesar (4 specimens), Safi Abad (3 specimens), 5-16 May 2010 (3 specimens), 5-16 May 2011 (4 specimens), *leg.* Elahe Shojaei.

**Flowers visited:** White umbellifers; *Caltha* sp., *Convolvulus* sp., *Euphorbia* sp., *Prunus padus*, *Ranunculus* sp., *Salix repens*, *Taraxacum* sp., *Tussilago* sp., *Ulex* sp. (Speight, 2011).

**Distribution in Iran:** Gilan, Isphahan, Maraghe, Semnan, Tehran (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Vosughian *et al.*, 2013)

# Eupeodes luniger Meigen, 1822

**Material examined:** 2 specimens ( $2 \stackrel{\frown}{\downarrow} \stackrel{\frown}{\downarrow}$ ), Robat Sang (2 specimens), 5-16 May 2011 (2 specimens), *leg.* Elahe Shojaei.

**Flowers visited:** White Umbelliferae; Calluna sp., Leontodon sp., Malus sylvestris, Polygonum cuspidatum, Prunus spinosa, Ranunculus sp., Rosa rugosa, Senecio sp., Taraxacum sp. (Speight, 2011).

**Distribution in Iran:** Fars, Golestan, Maraghe, Mazandaran, Sistan and Baluchistan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Vosughian *et al.*, 2013).

#### Eupeodes nuba Wiedemann, 1830

**Distribution in Iran:** Ardebil, East Azerbayjan, Fars, Gorgan, Gilan, Ilam, Khuzestan, Razavi Khorasan (Kashmar, Mashhad and Neyshabor), Semnan, Sistan and Baluchistan, Tehran (Sadeghi, Kayvanfar and Jajvandan, 2006; Gilasian, 2007; Ehteshamnia *et al.*, 2010; Jalilian *et al.*, 2010).

#### Ischiodon (Simosyrphus) scutellaris Fabricius, 1805

**Material examined:** 10 specimens ( $3 \circlearrowleft \circlearrowleft$ ,  $7 \circlearrowleft \circlearrowleft$ ), Robat Sang (2 specimens), Safi Abad (8 specimens), 5-16 May 2010 (7specimens), 5-16 May 2011 (3specimens), leg. Elahe Shojaei.

Flowers visited: Citrullus colocynthis, Diplotaxis harra, Fagonia indica, Reichardia sp. (Speight, 2011).

**Distribution in Iran:** Fars, Hormozgan, Kerman, Khuzestan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007).

#### Melanostoma mellinum Linnaeus, 1758

**Material examined:** 4 specimens (13, 399), Hesar (4 specimens), 4-16 September 2010 (2 specimens), 5-16 May 2011 (2 specimens), leg. Elahe Shojaei.

**Flowers visited:** Graminae; Cyperaceae; white Umbelliferae; *Allium ursinum*, *Bellis perennis*, *Caltha* sp., *Eschscholzia californica*, *Euphorbia* sp., *Leontodon* sp., *Luzula* sp., *Plantago* sp., *Ranunculus* sp., *Salix repens*, *Stellaria holostea*, *Succisa* sp., *Taraxacum* sp. (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Gilan, Gorgan, Ilam, Khuzestan, Maraghe, Mazandaran (Sadeghi, Kayvanfar & Jajvandan, 2006; Jalilian *et al.*, 2010; Vosughian *et al.*, 2013).

# Paragus albifrons Fallen, 1817

**Material examined:** 8 specimens ( $3 \circlearrowleft \circlearrowleft$ ,  $5 \hookrightarrow \hookrightarrow$ ), Hesar (4 specimens), Robat Sang (3 specimens), Safi Abad (1 specimen), 11-22 July 2010 (3 specimens), 4-16 September 2010 (5 specimens), *leg.* Elahe Shojaei.

Flowers visited: umbellifers; Ranunculus sp., Sambucus ebulus, Veronica sp. (Speight, 2011).

**Distribution in Iran:** East Azerbayjan, Ilam, Tehran (Ehteshamnia et al., 2010; Jalilian et al., 2010; Ahmadian & Pashaei Rad, 2012).

#### Paragus bicolor Fabricius, 1794

**Material examined:** 20 specimens ( $5 \circlearrowleft \circlearrowleft$ ,  $15 \circlearrowleft \circlearrowleft$ ), Hesar (3 specimens), Robat Sang (9 specimens), Safi Abad (8 specimens), 5-16 May 2010 (4 specimens), 11-22 July 2010 (5 specimens), 4-16 September 2010 (7 specimens), 5-16 May 2011 (4 specimens), leg. Elahe Shojaei.

**Flowers visited:** Euphorbia sp., Herniaria glabra, Potentilla sp., Sedum sp., Scleranthus sp., Solidago sp. (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Gilan, Gorgan, Ilam, Khuzestan, Kordestan, Maraghe, North Khorasan, Razavi Khorasan (Kashmar, Mashhad and Neyshabor), Semnan, West Azerbayjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Ehteshamnia *et al.*, 2010; Jalilian *et al.*, 2010; Khaghaninia and Hosseini, 2013; Vosughian *et al.*, 2013).

### Paragus haemorrhous Meigen, 1822

**Material examined:** 2 specimens  $(2 \stackrel{\frown}{\hookrightarrow} \stackrel{\frown}{\hookrightarrow})$ , Hesar (1specimen), Safi Abad (1 specimen), 11-22 July 2010 (2 specimens), *leg.* Elahe Shojaei.

Flowers visited: umbellifers; Calluna sp., Jasione montana, Matricaria sp., Origanum sp.,

Polygonum sp., Potentilla anserina, P. erecta, P. fruticosa, Solidago sp., Stellaria sp. (Speight, 2011).

**Distribution in Iran:** East Azerbayjan, Fars, Gilan, Kordestan, Maraghe, Razavi Khorasan (Mashhad) (Sadeghi, Kayvanfar & Jajvandan, 2006; Ehteshamnia *et al.*, 2010; Khaghaninia & Hosseini, 2013; Vosughian *et al.*, 2013)

#### Paragus quadrifasciatus Meigen, 1822

**Material examined:** 3 specimens ( $3 \stackrel{\frown}{\hookrightarrow} \stackrel{\frown}{\hookrightarrow}$ ), Robat Sang (2 specimens), Safi Abad (1 specimen), 11-22 July 2010 (2 specimens), 4-16 September 2010 (1 specimens), leg. Elahe Shojaei.

Flowers visited: yellow composites (Speight, 2011).

**Distribution in Iran:** East Azerbayjan, Ilam, Kordestan, Razavi Khorasan (Mashhad and Neyshabor), Semnan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Jalilian *et al.*, 2010).

# Paragus tibialis Fallen, 1817

**Material examined:** 3 specimens ( $3 \stackrel{\frown}{\hookrightarrow} \stackrel{\frown}{\hookrightarrow}$ ), Robat Sang (3 specimens), 5-16 May 2010 (1 specimen), 4-16 September 2010 (2 specimens), *leg.* Elahe Shojaei.

Flowers visited: Jasione montana, Potentilla sp., Salix repens (Speight, 2011).

**Distribution in Iran:** East Azerbayjan, Kordestan, Maraghe, Mazandaran, Tehran, Zanjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Khaghaninia & Hosseini, 2013; Vosughian *et al.*, 2013).

#### Scaeva albomaculata Macquart, 1842

**Material examined:** 2 specimens ( $2 \stackrel{\frown}{\downarrow} \stackrel{\frown}{\downarrow}$ ), Hesar (2 specimens), 5-16 May 2010 (2 specimens), *leg.* Elahe Shojaei.

Flowers visited: Cytisus sp., Senecio sp., Thapsia sp. (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Gorgan, Ilam, Isfahan, Kerman, Khuzestan, Razavi Khorasan (Mashhad), Tehran (Gilasian, 2007; Sadeghi, Kayvanfar & Jajvandan, 2006; Ehteshamnia *et al.*, 2010; Jalilian *et al.*, 2010).

#### Scaeva latimaculata Brunetti, 1923

**Material examined:** 2 specimens ( $1 \circlearrowleft$ ,  $1 \circlearrowleft$ ), Robat Sang (2 specimens), 4-16 September 2010 (2 specimens), *leg.* Elahe Shojaei.

Flowers visited: yellow composites (Speight, 2011).

**Distribution in Iran:** East Azerbayjan, Gorgan, Ilam, Kerman, Razavi Khorasan (Kashmar), Sistan and Baluchistan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Jalilian *et al.*, 2010).

# Sphaerophoria rueppelli Wiedemann, 1830

**Material examined:** 23 specimens ( $8 \circlearrowleft \circlearrowleft , 15 \hookrightarrow \circlearrowleft$ ), Hesar (10 specimens), Robat Sang (9 specimens), Safi Abad (4 specimens), 5-16 May 2010 (4 specimens), 11-22 July 2010 (3 specimens), 4-16 September 2010 (7 specimens), 5-16 May 2011 (9 specimens), leg. Elahe Shojaei.

**Flowers visited:** yellow composites; Asparagus sp., Aster sp., Limonium sp., Matricaria sp., Polygonum sp., Ranunculus sp., Rumex sp., Sinapis sp., Tamarix sp. (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Fars, Gorgan, Ilam, Khuzestan, Mazandaran, Razavi Khorasan (Kashmar and Mashhad), Semnan, Sistan and Baluchistan, West Azerbayjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Ehteshamnia *et al.*, 2010; Jalilian *et al.*, 2010).

# Sphaerophoria scripta Linnaeus, 1758

**Material examined:** 58 specimens ( $21 \circlearrowleft \circlearrowleft$ ,  $37 \circlearrowleft \circlearrowleft$ ), Hesar (30 specimens), Robat Sang (11 specimens), Safi Abad (17 specimens), 5-16 May 2010 (19 specimens), 11-22 July 2010 (4 specimens), 4-16 September 2010 (10 specimens), 5-16 May 2011 (25 specimens), leg. Elahe Shojaei.

Flowers visited: white umbellifers; Achillea sp., Campanula rapunculoides, Cirsium arvense, Crataegus sp., Erigeron sp., Eschscholzia californica, Euphorbia sp., Leontodon sp., Origanum vulgare, Prunus spinosa, Ranunculus sp., Tripleurospermum inodoratum, Tussilago sp. (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Fars, Gilan, Golestan, Gorgan, Hamedan, Ilam, Isphahan, Khuzestan, Maraghe, Mazandaran, North Khorasan, Qazvin, Razavi Khorasan (Kashmar, Mashhad and Neyshabor), Sistan and Baluchistan, West Azerbayjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Jalilian *et al.*, 2010; Vosughian *et al.*, 2013).

# Sphaerophoria turkmenica Bankowska, 1964

**Material examined:** 47 specimens  $(17 \circlearrowleft \circlearrowleft, 30 \circlearrowleft \circlearrowleft)$ , Hesar (31 specimens), Robat Sang (9 specimens), Safi Abad (7 specimens), 5-16 May 2010 (14 specimens), 11-22 July 2010 (4 specimens), 4-16 September 2010 (6 specimens), 5-16 May 2011 (23 specimens), leg. Elahe Shojaei.

Flowers visited: no data (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Fars, Gorgan, Ilam, Isphahan, Khuzestan, Mazandaran, North Khorasan, West Azerbayjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Jalilian *et al.*, 2010).

# Syrphus ribesii Linnaeus, 1758

**Material examined:** 1 specimen ( $1^{\circ}$ ), Robat Sang (1 specimen), 5-16 May 2010 (1 specimens), *leg.* Elahe Shojaei.

Flowers visited: visits a wide range of yellow, white, pink and blue flowers, including composites and umbellifers and the flowers of many trees and shrubs (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Fars, Gorgan, Hamedan, Isfahan, Maraghe, Mazandaran, Razavi Khorasan (Kashmar and Mashhad), Sistan and Baluchistan, Tehran (Sadeghi, Kayvanfar & Jajvandan, 2006; Gilasian, 2007; Vosughian *et al.*, 2013).

# Subfamily Eristalinae

# Eristalinus aeneus Scopoli, 1763

**Material examined:** 6 specimens  $(2 \circlearrowleft \circlearrowleft, 4 \hookrightarrow \circlearrowleft)$ , Hesar (2 specimens), Robat Sang (1 specimen), Safi Abad (3 specimens), 5-16 May 2010 (2 specimens), 5-16 May 2011 (4 specimens), *leg.* Elahe Shojaei. **Flowers visited:** yellow composites, white umbellifers, *Aster* sp., *Berteroa incana*, *Cistus* sp., *Origanum* sp., *Salix repens*, *Taraxacum* sp. (Speight, 2011).

**Distribution in Iran:** East Azerbayjan, Gilan, Gorgan, Hamedan, Ilam, Khuzestan, North Khorasan, Razavi Khorasan (Kashmar, Mashhad and Neyshabor), Semnan, West Azerbayjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Jalilian *et al.*, 2010).

#### Eristalinus taenoips Wiedemann, 1818

Material examined: 1 specimen (1♀), Hesar (1 specimen), 5-16 May 2011 (1specimen), leg. Elahe Shojaei.

**Flowers visited:** umbellifers; Calluna vulgaris, Cistus sp., Daphne sp., Eryngium sp., Hedera sp., Mentha sp., Rubus sp., Senecio sp., Solidago sp. (Speight, 2011).

**Distribution in Iran:** East Azerbayjan, Gorgan, Khuzestan, Ilam (Sadeghi, Kayvanfar & Jajvandan, 2006; Jalilian *et al.*, 2010)

# Eristalis arbustorum Linnaeus, 1758

**Material examined:** 12 specimens ( $4 \circlearrowleft \circlearrowleft$ ,  $8 \hookrightarrow \circlearrowleft$ ), Hesar (3 specimens), Robat Sang (5 specimens), Safi Abad (4 specimens), 5-16 May 2010 (3 specimens), 4-16 September 2010 (2 specimens), 5-16 May 2011 (7 specimens), *leg.* Elahe Shojaei.

**Flowers visited:** visits the flowers of a wide range of low-growing plants and shrubs (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Gilan, Gorgan, Hamedan, Ilam, Isphahan, Khuzestan, Mazandaran, Razavi Khorasan (Mashhad and Neyshabor), Semnan, Sistan and Baluchistan, West Azerbayjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Jalilian *et al.*, 2010).

### Eristalis tenax Linnaeus, 1758

**Material examined:** 2 specimens ( $2 \stackrel{\frown}{\hookrightarrow} \stackrel{\frown}{\hookrightarrow}$ ), Safi Abad (2 specimens), 5-16 May 2011 (2 specimens), *leg.* Elahe Shojaei.

**Flowers visited:** visits a wide range of flowers, including white, yellow, pink and blue blooms (Speight, 2011).

**Distribution in Iran:** Ardebil, East Azerbayjan, Gilan, Gorgan, Hamedan, Ilam, Khuzestan, Mazandaran, Razavi Khorasan (Mashhad and Neyshabor), Semnan, West Azerbayjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Jalilian *et al.*, 2010).

# Eumerus sogdianus Stackelberg, 1952

**Material examined:** 5 specimens (13, 499), Hesar (2 specimens), Robat Sang (1 specimen), Safi Abad (2 specimens), 11-22 July 2010 (2 specimens), 4-16 September 2010 (2 specimens), 5-16 May 2011 (1 specimens), *leg.* Elahe Shojaei.

Flowers visited: white umbellifers; Sedum acre (Speight, 2011).

**Distribution in Iran:** Ilam, West Azerbayjan (Sadeghi, Kayvanfar & Jajvandan, 2006; Jalilian *et al.*, 2010).

# Eumerus strigatus Fallen, 1817

**Material examined:** 3 specimens ( $3 \stackrel{\frown}{\hookrightarrow} \stackrel{\frown}{\hookrightarrow}$ ), Hesar (2 specimens), Robat Sang (1 specimen), 11-22 July 2010 (1 specimen), 4-16 September 2010 (2 specimens), *leg.* Elahe Shojaei.

**Flowers visited:** umbellifers; *Allium ursinum*, *Convolvulus* sp., *Eschscholzia californica*, *Euphorbia* sp., *Fragaria* sp., *Leontodon* sp., *Papaver* sp., *Potentilla erecta*, *Ranunculus* sp., *Sonchus arvensis*, *Taraxacum* sp. (Speight, 2011).

Distribution in Iran: Kerman, Tehran (Sadeghi, Kayvanfar & Jajvandan, 2006).

# Eumerus tuberculatus Rondani, 1857

**Material examined:** 5 specimens ( $2 \circlearrowleft \circlearrowleft$ ,  $3 \hookrightarrow \hookrightarrow$ ), Hesar (2 specimens), Robat Sang (3 specimens), 5-16 May 2010 (1 specimens), 11-22 July 2010 (2 specimens), 4-16 September 2010 (2 specimens), *leg.* Elahe Shojaei.

Flowers visited: Euphorbia sp., Fragaria sp., Leucanthemum sp., Ranunculus sp. (Speight, 2011).

Distribution in Iran: Gorgan (Sadeghi, Kayvanfar & Jajvandan, 2006).

#### Neoascia oblique Coe, 1940

**Material examined:** 1 specimen ( $1^{\circ}$ ), Hesar (1 specimen), 5-16 May 2011(1 specimen), *leg.* Elahe Shojaei.

Flowers visited: white umbellifers; Galium sp., Ranunculus sp., Taraxacum sp. (Speight, 2011).

Distribution in Iran: East Azerbaijan, Kordestan (Khaghaninia & Hosseini, 2013).

# Syritta pipiens Linnaeus, 1758

Material examined: 56 specimens  $(22 \circlearrowleft \circlearrowleft \circlearrowleft 34 \circlearrowleft \circlearrowleft)$ , Hesar (25 specimens), Robat Sang (17specimens), Safi Abad (14 specimens), 5-16 May 2010 (16 specimens), 11-22 July 2010 (11specimens), 4-16 September 2010 (13specimens), 5-16 May 2011 (16 specimens), leg. Elahe Shojaei.

Flowers visited: white umbellifers; Achillea sp., Allium sp., Aster sp., Calluna sp., Cardamine sp., Cirsium palustre, Convolvulus sp., Crataegus sp., Epilobium sp., Euphorbia sp., Galium sp., Jasione montana, Leontodon sp., Polygonum cuspidatum, Potentilla erecta, Prunus laurocerasus, Ranunculus sp., Rosa canina, Senecio jacobaea, Sorbus aucuparia, Tussilago sp. (Speight, 2011).

**Distribution in Iran:** Ardebil, Gilan, Gorgan, Ilam, Isphahan, Khuzestan, Mazandaran, Razavi Khorasan (Mashhad and Neyshabor), Semnan (Sadeghi, Kayvanfar & Jajvandan, 2006; Jalilian *et al.*, 2010).

#### **ACKNOWLEDGMENTS**

We are grateful to Dr. Anatoli Barkalov (Siberian Zoological Museum, Novosibirsk, Russia) for his help with the species identification and sending some papers about syrphid flies.

#### LITERATURE CITED

Ahmadian, S.A., Pashaei Rad, Sh., 2012. Four species *Metasyrphus latilunulatus*, *Chrysogaster hirtlla*, *Merodon elegans* and *Eumerus ornatus* (Diptera: Syrphidae) newly reported from Iran. *Journal of Animal Environment* 4(1), 79-84.

Ahmadian, S.A., Pashaei Rad, Sh., 2012. Fauna and Diversity of hoverflies (Diptera: Syrphidae) in Damavand. *Journal of Animal Environment* 4, 49-64.

Amiri Moghadam, F., 2004. Survey of hoverflies (Diptera: Syrphidae) in Kashmar and Bardaskan [in Persian with English summary]. M.Sc. Dissertation, Department of Biology, Faculty of Science, Shahid Beheshti University, Tehran, p. 141.

Ashrafi, F., Pashaei Rad, Sh., 2010. A new record of the subfamily Syrphinae (Diptera: Syrphidae) for the Iranian fauna. *Zoology in the Middle East* 51, 119-120.

Barkalov, A., 2013. A new Platycheirus Le Peletier et Serville, 1828 (Diptera, Syrphidae) species of the manicatus subgroup, from the Taimyr Peninsula (Northern Siberia). *Zootaxa* 3681(2), 175–181.

Bei- Bienko, G., 1988. Keys to the insects of the European part of the USSR (Volume V) Diptera and Syphonaptera. Brill Archive. Part II. Smithsonian Institution Libraries and the National Science Foundation Washington, D.C. 10, p. 148.

Dousti, A.F., 1999. Fauna and Diversity of Syrphid flies in Ahwaz region. Master Thesis, Shahid-Chamran University, p. 129.

Dousti, A.F., Hayat, A., 2006. A catalogue of the Syrphidae (Insecta: Diptera) of Iran. *Journal of the Entomological Research Society* 8(3), 5-38.

Ehteshamnia, N., Khaghaninia, S., Farshbaf Pourabad, R., 2010. Some hoverflies of subfamily Syrphinae of Qurigol fauna in East Azerbayjan province, Iran (Diptera: Syrphidae). *Munis Entomology and Zoology journal* 5(2), 499-505.

Ghahari, H., Hayat, R., Tabari, M., Ostovan, H., 2008. Hoverflies (Diptera: Syrphidae) from rice fields and around grasslands of northern Iran. *Munis Entomology and Zoology journal* 3(1), 275-284.

Gharali, B., Alichi, M., Radjabi, Gh., 2000. The new records of syrphid flies (Diptera: Syrphidae). Proceedings of the 14<sup>th</sup> Iranian Plant Protection Congress, p. 348.

Gilasian, E., 2007. Study on Tribe Syrphini in Iran. Journal of Entomological Society of Iran 27(1), 86-107.

Gilasian, E., 2005. New record of one genus and six species of Syrphidae (Diptera) from Iran. *Journal of Entomological Society of Iran* 25(1), 75-76.

Gilasian, E., Sorokina, V.S., 2011. The genus Paragus Latreille (Diptera: Syrphidae) in Iran, with the description of a new species. *Zootaxa* 2764, 49–60.

Gilbert, F.S., 1981. Foraging ecology of hoverflies: Morphology of the mouthparts in relation to feeding on nectar and pollen in some common urban species. *Ecological Entomology* 6, 245-262.

Hosseini, M., Sadeghi, H., 2007. Diversity of Syrphidae (Diptera) in agroecosystems of Neyshabor region. *Plant Protection Journal* 22(2), 85-93.

Jalilian, F., 2014. Evaluation of population fluctuation of predacious species of syrphidflies (Dip: Syrphidae) on Cabbage aphid *Brevicoryne brassicae* in rapeseed fields. *Agricultural Pest Management* 1(1), 46-54.

Jalilian, F., Fathipour, Y., Talebi, A.A., edaratian, A., 2010. Faunal and Population Studies of Syrphidflies (Diptera: Syrphidae) in Ilam. *Environmental Sciences* 7(4), 73-86.

Kazerani, F., Talebi, A.A., Gilasian, E., Steenis, J.V., 2012. New record of *Spilomyia triangulate* Van Steenis, 2000 (Diptera: syrphidae) from Iran. *Acta entomolica serbica* 17(1/2), 39-44.

Kazerani, F., Talebi, A.A., Gilasian, E., Khayrandish, M., 2014. New record of the genus and species *Temnostoma vespiforme* (Diptera: Syrphidae) from Iran. *Journal of Crop Protection* 3(1), 69-73.

Khaghaninia, S., 2010. Faunistic study on flower flies of Zunuz region in East Azerbaijan province-Iran. *Munis Entomology and Zoology journal* 5(2), 586-93.

Khaghaninia, S., Hosseini, C., 2013.Taxonomic study of Paragus Latreille (Diptera: Syrphidae) in the East Azerbayjan and Kordestan provinces of Iran. *Efflatounia* 13, 8-18.

Kayvanfar, N., Bagherian, A., 2006. A Morphological Study of Syrphidae in Mashhad. *Iranian Journal of Biology* 18(4), 332-340.

Mehrabi, R., Pashaei Rad, Sh., Shiravi, A.H., Amiri Moghaddam, F., 2006. Determination of the Biodiversity and Fauna of Hoverflies (Dip: Syrphidae) in Damghan (Semnan province) - Introducing 23 New Records from Semnan Province. *Environmental Sciences* 13(1), 61-68.

Nami, M.H., 2010. National gazetteer of the Islamic Republic of Iran (province of Razavi Khorasan: Khaf township). National Geographical Organization publication. Tehran (in Persian). pp 161.

Rotheray, G., Gilbert, F., 1999. Phylogeny of Palearctic Syrphidae (Diptera) Evidence from larval stages. Zoological Journal of the Linnean Society 127, 1-112.

Sadeghi, H., Kayvanfar, N., Jajvandian, R., 2006. An introduction to Syrphidae. Ferdowsi University of Mashhad Publication (In Persian). Sonbole press. p. 236.

Sommaggio, D., 1999. Syrphidae: can they be used as an environmental bioindicators? *Agriculture, Ecosystems and Environment* 74(3), 343-356.

Sorokina, V.S., 2009. Hoverflies of the genus Paragus Latr. (Diptera, Syrphidae) of Russia and adjacent countries. *Entomological Review* 89(3), 351-366.

Speight, M.C.D., 2011. Species accounts of European Syrphidae (Diptera), Glasgow 2011. Syrph the Net, the database of European Syrphidae 65, p. 292.

Speight, M.C.D., 2010. Species accounts of European Syrphidae (Diptera) 2010. Syrph the Net, the database of European Syrphidae 59, p. 285.

Stubbs, A.E., Falk, S.J., 2002. British hover flies. An Illustrated Identification Guide. The British Entomology and Natural History Society, p. 469.

Thompson, F.C., Rotheray, G.E., 1998. Family Syrphidae. In: Papp, L., Darvas, B. (Eds.), Contributions to a Manual of Palaearctic Diptera (with special reference to flies of economic importance). Vol. 3, Science Herald, Budapest, pp. 81-139.

Vockeroth, J.R., Tompson, F.C., 1981. Syrphidae in: Manual of Nearctic Diptera. Biosystematic Research Institute. Ottawa-Ontario 2, p. 674.

Vosughian, M., Khaghaninia, S., Haddad Irani Nejad, K., 2013. Hoverfly fauna of the subfamily Syrphinae (Diptera: Syrphidae) of Maragheh region, Iran, with a new record for the Iranian insect fauna. *Calodema* 274, 1-6.